

# Lower CS-Closed Sets and Functions

Charki Amara and Marc Ciligot-Travain

Département de Mathématiques Université d'Avignon 33 rue Pasteur

metadata, citation and similar papers at [core.ac.uk](http://core.ac.uk)

Submitted by Muhammad Imdad, 11/04/1999

Received March 12, 1999

In this work, we introduce a class of convex sets,  $LCSC^{\mathcal{F}}(X)$ , of a locally convex separated and *not necessarily separable* topological vector space  $X$ . They are called the lower CS-closed sets. This class contains the CS-closed sets, satisfies the property  $\text{core}(C) = \text{int}(C)$ ,  $\forall C \in LCSC^{\mathcal{F}}(X)$  when  $X$  is metrizable barrelled, and is stable under many operations. Among them, the projection and the denumerable intersection. We characterize the lower CS-closed functions (i.e., the functions who have a lower CS-closed epigraph) as marginal functions of CS-closed ones and show that they are very stable too. We establish an open mapping and a closed graph theorem for the lower CS-closed relations. Finally, we show that every real extended valued lower CS-closed function defined on a metrizable barrelled space is continuous on the interior of its domain. This result allows us to extend classical theorems of convex duality by replacing lower semicontinuous functions by lower CS-closed ones. More than that, it systematizes and extends some methods of convex analysis. © 1999 Academic Press

*Key Words:* convex analysis; CS-closed; duality; openness.

## 1. INTRODUCTION

Consider a convex set  $C$  of a locally convex Hausdorff (separated) topological vector space  $X$  (a l.c.s. space in the rest of the paper). We are interested in the property

$$\text{core}(C) = \text{int}(C), \quad (1)$$

where  $\text{core}(C)$  is the algebraic interior of  $C$ , defined as

$$\text{core}(C) = \{x \in C \mid \forall u \in X, \exists \varepsilon > 0, x + \varepsilon[-u, u] \subset C\},$$

and  $\text{int}(C)$  denotes the interior of  $C$ .

The reason why we are interested in the equality (1) is that this property (possibly used with the Hahn–Banach theorem) has many applications in convex analysis: continuity of convex functions, open mapping theorem, closed graph theorem, calculus of conjugates, subdifferential calculus, etc.

One always has  $\text{int}(C) \subset \text{core}(C)$ . If  $\text{int}(C) \neq \emptyset$  or  $\text{core}(C) = \emptyset$  then (1) holds; but it is well known that (1) fails in general. For example, take for  $X$  the space of all continuous real functions defined on a compact topological space  $K$  (with  $K$  infinite) endowed with the pointwise convergence topology, and consider the set  $C = \{f \in X \mid \forall t \in K, f(t) > 0\}$ ; thus  $\text{core}(C) = C$  but  $\text{int}(C) = \emptyset$ .

In fact to ensure that the relation (1) holds, one needs some conditions on  $C$  and/or on the whole space  $X$ .

If  $X$  is a finite dimensional, any convex set of  $X$  satisfies (1). Any closed convex set in a barrelled space satisfies (1). A larger class of convex subsets satisfying (1) is given by the CS-closed sets (whose definition is recalled below) in a metrizable barrelled space.

All the classes of sets considered above have one same default; they are not sufficiently stable: essentially, the projection of a closed convex (resp. a CS-closed set) is not necessarily closed (resp. CS-closed).

One approach to define a class of convex sets sufficiently stable and containing the closed convex sets is to add measurability assumptions. One can consult Kusraev [12] and references therein. But it seems that it needs the space  $X$  to be *separable* (or to introduce a similar hypothesis).

The purpose of this paper is to introduce a class of convex sets, containing the CS-closed sets, namely the lower CS-closed sets, which satisfies (1), enjoys remarkable stability properties, and does not need the space  $X$  to be separable.

In fact a lower CS-closed set of a l.c.s. space  $X$  is just the projection on  $X$  of a CS-closed set of a product space  $X \times Y$  with  $Y$  Fréchet.

This class of sets appears implicitly in former works. However, it seems that it has never been studied (neither defined!). One can see the class of lower CS-closed sets as the result of the systematization (and extension) of techniques used before in convex analysis concerning the problems evoked above.

We will see that any lower CS-closed set of a metrizable barrelled space satisfies (1).

As a direct consequence of the definition, the projection of a lower CS-closed set of a Fréchet space is lower CS-closed too. Moreover the sum of two lower CS-closed sets is always lower CS-closed. In particular the sum of two closed linear spaces is always lower CS-closed but may fail to be CS-closed.

Here we develop some applications concerning the continuity and the subdifferentiability of a lower CS-closed function (a function whose epigraph is lower CS-closed).

We observe that in practice many convex functions are lower CS-closed but not lower semicontinuous or CS-closed. Below, we prove that the sum, the max, the infimal convolution, the level sum, and the composite of two lower CS-closed functions are lower CS-closed. We characterize a lower CS-closed function as a marginal function of a CS-closed one.

As an application, we give a general duality result involving lower CS-closed perturbation functional. In this way, we obtain also some calculus rules about asymptotic functionals under mild assumptions.

## 2. LOWER CS-CLOSED SETS AND FUNCTIONS

### 2.1. Preliminaries: Some Properties of CS-Closed Sets

Let us first recall the definition of the CS-closed sets. Let  $C$  be a subset of a l.c.s. space. By a convex series of elements of  $C$ , we mean a series of the form  $\sum_{n \in \mathbb{N}} \lambda_n x_n$ , where  $(\forall n \in \mathbb{N}, x_n \in C)$ ,  $(\forall n \in \mathbb{N}, \lambda_n \geq 0)$ , and  $\sum_{n=0}^{+\infty} \lambda_n = 1$ . We say that  $C$  is CS-closed (some authors say ideally convex) if it contains the sum of every convergent convex series of its elements. Note that any closed convex set is CS-closed. The CS-closed sets are always convex but not necessarily closed: for example, every open convex set is CS-closed. In the sequel we shall use some properties of the CS-closed sets we recall below.

( $\mathcal{N}_1$ ) If  $\{X_i\}_{i \in I}$  is a family of l.c.s. spaces and for each  $i \in I$ ,  $A_i \subset X_i$  is a CS-closed set then  $\prod_{i \in I} A_i$  is a CS-closed set of  $\prod_{i \in I} X_i$  endowed with the product topology.

( $\mathcal{N}_2$ ) Every intersection of CS-closed sets is CS-closed.

If  $A$  is a CS-closed set of a l.c.s. space  $X$  and  $M$  is a subspace of  $X$  then  $A \cap M$  is a CS-closed set of  $M$  endowed with the induced topology.

One says that a function  $f: X \rightarrow \overline{\mathbb{R}}$  is CS-closed if its epigraph,  $\text{epi } f$ , defined as  $\text{epi } f = \{(x, r) \in X \times \mathbb{R}, f(x) \leq r\}$ , is a CS-closed subset of  $X \times \mathbb{R}$ .

Due to ( $\mathcal{N}_1$ ) and ( $\mathcal{N}_2$ ), the indicator function of a subset  $A$  of  $X$ , that is, the function  $\delta_A$  defined by  $\delta_A(x) = 0$  if  $x \in A$  and  $\delta_A(x) = +\infty$  if  $x \notin A$ , is a CS-closed function if and only if  $A$  is CS-closed.

Note that every convex lower semicontinuous function is CS-closed.

One can refer to Jameson [11], Lifshits [15], Holmes [10], or Kusraev and Kutateladze [13] for further details about the CS-closed sets.

### 2.2. Lower CS-Closed Sets and Functions

Let  $A$  be a subset of a product space  $X \times Y$ ; we denote by  $A_X$  the projection of  $A$  on the space  $X$ . Recall that a Fréchet space is a locally convex topological vector space which is metrizable and complete.

Let us introduce now the classes of sets and functions we are interested in.

**DEFINITION 2.1.** A subset  $C$  of a l.c.s. space  $X$  is said to be lower CS-closed if there exists a Fréchet space  $Y$  and a CS-closed set  $A$  of  $X \times Y$  such that  $C = A_X$ .

We denote by  $LCSC^{\mathcal{F}}(X)$  the set of all lower CS-closed subsets of  $X$ .

Every CS-closed set is a lower CS-closed set but the converse is false: for example, we will see that the sum of two lower CS-closed sets is always lower CS-closed but may fail to be CS-closed.

Let us give an example of lower CS-closed set: take a CS-closed function  $f: X \rightarrow \mathbb{R}$ ; the domain of  $f$ ,  $\text{dom } f = \{x \in X, f(x) < +\infty\}$ , is a lower CS-closed set because it is the projection of the epigraph of  $f$  ( $\text{dom } f = [\text{epi } f]_X$ ).

We extend Definition 2.1 to functions and give a link between lower CS-closed sets and lower CS-closed functions.

**DEFINITION 2.2.** A function  $f: X \rightarrow \overline{\mathbb{R}}$  defined on a l.c.s. space  $X$  is said to be lower CS-closed if its epigraph is a lower CS-closed subset of  $X \times \mathbb{R}$ .

**LEMMA 2.1.** A subset  $C$  of a l.c.s. space  $X$  is lower CS-closed if and only if  $\delta_C$  is a lower CS-closed function.

### 2.3. Stability of Lower CS-Closed Sets

The main motivation for introducing this class of sets and functions lies in its simplicity and its strong stability regarding various operations.

**THEOREM 2.1.** Let  $C$  be a lower CS-closed set of a product  $X \times Y$  of l.c.s. spaces where  $Y$  is a Fréchet space. Then  $C_X$  is a lower CS-closed set.

Another interesting fact is:

**THEOREM 2.2.** Lower CS-closed sets still satisfy  $(\mathcal{H}_1)$  and  $(\mathcal{H}_2)$  for denumerable families.

*Proof.* We prove only that the intersection of a denumerable family of lower CS-closed sets is lower CS-closed.

Let  $\{C_n\}_{n \in \mathbb{N}}$  be a family of lower CS-closed sets of a l.c.s. space  $X$ . For each  $n \in \mathbb{N}$ , there exists a Fréchet space  $Y_n$  and a CS-closed subset  $A_n$  of  $X \times Y_n$  such that  $C_n = [A_n]_X$ . Set  $Y = \prod_{n \in \mathbb{N}} Y_n$  and  $\hat{A}_n = \{(x, (y_k)_{k \in \mathbb{N}}) \in X \times Y \mid (x, y_n) \in A_n\}$ . Then  $Y$  is a Fréchet space and  $\hat{A}_n$  is CS-closed. So  $\bigcap_{n \in \mathbb{N}} \hat{A}_n$  is CS-closed too. Using the fact that

$$\bigcap_{n \in \mathbb{N}} C_n = \left[ \bigcap_{n \in \mathbb{N}} \hat{A}_n \right]_X,$$

we obtain the result. ■

One can deduce from these properties various corollaries. We identify a set-valued mapping with its graph.

**COROLLARY 2.1.** *If  $A$  is a lower CS-closed set of a Fréchet space  $X$  and  $T$  is a lower CS-closed set of  $X \times Y$  where  $Y$  is a l.c.s. space, then  $T(A)$  is a lower CS-closed set of  $Y$ .*

*Proof.* By definition, we have  $T(A) = [T \cap A \times Y]_Y$ . So by Theorems 2.2 and 2.1,  $T(A)$  is lower CS-closed. ■

**COROLLARY 2.2.** *Let  $X$  be a l.c.s. space. The level sets of a lower CS-closed function  $f: X \rightarrow \mathbb{R}$ , that are the subsets of  $X$*

$$[f \leq r] = \{x \in X \mid f(x) \leq r\}, \quad [f < r] = \{x \in X \mid f(x) < r\}, \quad r \in \mathbb{R},$$

*are lower CS-closed sets of  $X$ .*

*Proof.* For each  $r \in \mathbb{R}$ ,

$$[f \leq r] = [\text{epi } f \cap (X \times ]-\infty, r])_X,$$

$$[f < r] = [\text{epi } f \cap (X \times ]-\infty, r[)]_X.$$

■

It is known that a linear subspace of a metrizable space is CS-closed if and only if it is closed. Moreover, since the sum of two closed linear subspaces of a Banach space may fail to be closed (see [6]), the sum of two CS-closed sets does not need to be CS-closed.

**COROLLARY 2.3.** *The sum of two lower CS-closed subsets of a Fréchet space is still lower CS-closed.*

*Proof.* The set  $A + B$  is the image of the lower CS-closed subset  $A \times B$  by the addition whose graph is a closed linear space. ■

**COROLLARY 2.4.** *Let  $M: X \rightarrow Y$  and  $N: Y \rightarrow Z$  be two set-valued maps where  $X, Y, Z$  are l.c.s. spaces with  $Y$  Fréchet. If  $M$  and  $N$  are lower CS-closed, then  $N \circ M$  is lower CS-closed.*

*Proof.* By definition, one has  $N \circ M = [M \times Z \cap X \times N]_{X \times Z}$ . ■

Let us consider a last case. Let  $X$  and  $Z$  be two l.c.s. spaces,  $A \subset X \times X$  and  $B \subset X \times Z$ . Recall that the right partial sum of  $A$  and  $B$  is the set

$$A \dot{+} B = \{(x, z) \in X \times Z \mid \exists z_1 \in Z, \exists z_2 \in Z, (x, z_1) \in A, \\ (x, z_2) \in B, z = z_1 + z_2\}.$$

**COROLLARY 2.5.** *The right partial sum of two lower CS-closed sets  $A$  and  $B$  of a space  $X \times Z$  where  $X$  is a l.c.s. space and  $Z$  is a Fréchet space is still lower CS-closed.*

*Proof.* We set

$$T = \{(u, x, y, z) \in X \times Z \times Z \times Z, z = x + y\}$$

$$C = \{(u, x, y) \in X \times Z \times Z \mid (u, x) \in A, (u, y) \in B\}.$$

Then  $T$  is a closed linear subspace and  $C$  is lower CS-closed. Thus  $A \dot{+} B = [T \cap C \times Z]_{X \times Z}$  is lower CS-closed too. ■

## 2.4. Characterization of Lower CS-Closed Functions

In this section, we prove that a lower CS-closed function can be described as marginal of a CS-closed function. We first observe that a function is lower CS-closed if and only if its strict epigraph is lower CS-closed. Recall that the strict epigraph of a function  $f: X \rightarrow \overline{\mathbb{R}}$  is given by  $\text{epi}_s f = \{(x, r) \in X \times \mathbb{R} \mid f(x) < r\}$ .

**PROPOSITION 2.1.** *Let  $X$  be a l.c.s. space and  $f: X \rightarrow \overline{\mathbb{R}}$ . Then  $f$  is lower CS-closed if and only if its strict epigraph is lower CS-closed.*

*Proof.* We first observe that, for all  $\varepsilon > 0$ ,

$$\text{epi}_s f = \text{epi } f \dot{+} (X \times ]0, +\infty[), \quad (2)$$

$$\text{epi}_s(f - \varepsilon) = \text{epi}_s f \dot{+} (X \times [-\varepsilon, +\infty[). \quad (3)$$

Now if  $f$  is lower CS-closed then (2) and Corollary 2.5 tell us that  $\text{epi}_s f$  is lower CS-closed.

Conversely, if  $\text{epi}_s f$  is lower CS-closed then, using Corollary 2.5 once more, (3) tells us that  $\text{epi}_s(f - \varepsilon)$  is lower CS-closed for all  $\varepsilon > 0$ . On the other hand one has

$$\text{epi } f = \bigcap_{\varepsilon > 0, \varepsilon \in \mathbb{Q}} \text{epi}_s(f - \varepsilon).$$

Therefore  $\text{epi } f$  is lower CS-closed, which completes the proof. ■

We are now in position to characterize the class of lower CS-closed functions.

**THEOREM 2.3.** *Let  $X$  be a l.c.s. space. A function  $f: X \rightarrow \overline{\mathbb{R}}$  is lower CS-closed if and only if there exists a Fréchet space  $Y$  and a CS-closed function  $F: X \times Y \rightarrow \overline{\mathbb{R}}$  such that for all  $x \in X$ ,  $f(x) = \inf_{y \in Y} F(x, y)$ .*

*Proof.* If  $\text{epi } f$  is lower CS-closed then there exists a Fréchet space  $Y$  and a CS-closed set  $A$  of  $X \times \mathbb{R} \times Y$  such that  $\text{epi } f = A_{X \times \mathbb{R}}$ . We then have

$$f(x) = \inf_{r \in \mathbb{R}} r + \delta_{\text{epi } f}(x, r) = \inf_{(y, r) \in Y \times \mathbb{R}} r + \delta_A(x, r, y).$$

Setting  $F(x, (y, r)) = r + \delta_A(x, r, y)$ , we get a CS-closed function. Moreover  $f(x) = \inf_{(y, r) \in Y \times \mathbb{R}} F(x, (y, r))$  with  $Y \times \mathbb{R}$  Fréchet.

Assume that there exists a Fréchet space  $Y$  and a CS-closed function  $F: X \times Y \rightarrow \overline{\mathbb{R}}$  such that  $f(x) = \inf_{y \in Y} F(x, y)$ . Then

$$\text{epi}_s f = [\text{epi}_s F]_{X \times \mathbb{R}}. \quad (4)$$

Moreover  $Y$  is a Fréchet space and by Proposition 2.1,  $\text{epi}_s F$  is lower CS-closed. Now, using Theorem 2.1, (4) tells us that  $\text{epi}_s f$  is lower CS-closed. Finally, applying Proposition 2.1 once more, we get the announced result. ■

## 2.5. Stability of Lower CS-Closed Functions

Due to the preceding characterization, we have the following result:

**COROLLARY 2.6.** *Let  $F: X \times Z \rightarrow \overline{\mathbb{R}}$  be a lower CS-closed function where  $X$  is a l.c.s. space and  $Z$  is a Fréchet space. Then the marginal function  $f: X \rightarrow \overline{\mathbb{R}}$ , defined by  $f(x) = \inf_{z \in Z} F(x, z)$ , is lower CS-closed.*

It is known that the sum of two CS-closed functions having affine continuous minorants is CS-closed. The next proposition says, in particular, that the sum of two CS-closed functions is always lower CS-closed.

**COROLLARY 2.7.** *Let  $X$  be a l.c.s. space,  $f: X \rightarrow \overline{\mathbb{R}}$  and  $g: X \rightarrow \overline{\mathbb{R}}$  two lower CS-closed functions. Then  $\alpha f + \beta g$  is a lower CS-closed function for all  $\alpha, \beta > 0$ .*

*Proof.* Essentially due to the fact that  $\text{epi}(f + g) = \text{epi } f \dot{+} \text{epi } g$  (we make the convention  $(+\infty) + (-\infty) = (-\infty) + (+\infty) = +\infty$ ). ■

Concerning the supremum of a family of lower CS-closed functions, we have the following result:

**COROLLARY 2.8.** *Let  $(f_n)_{n \in \mathbb{N}}$  be a sequence of lower CS-closed functions defined on a l.c.s. space  $X$ . Then  $\sup_{n \in \mathbb{N}} f_n$  is a lower CS-closed function.*

*Proof.* One has

$$\text{epi}\left(\sup_{n \in \mathbb{N}} f_n\right) = \bigcap_{n \in \mathbb{N}} \text{epi } f_n.$$

Then by Theorem 2.2,  $\sup_{n \in \mathbb{N}} f_n$  is lower CS-closed. ■

The next result concerns the case of a lower CS-closed composite function:

**COROLLARY 2.9.** *Let  $X$  be a l.c.s. space, let  $Z$  be a Fréchet space preordered by a convex cone  $Z_+$ , let  $h$  be a mapping defined on a convex set  $\text{dom } h$  of  $X$  with values in  $Z$  which is convex with respect to  $Z_+$ , and let  $g: Z \rightarrow \overline{\mathbb{R}}$  be a convex function nondecreasing on  $h(\text{dom } h) + Z_+$  with respect to the preorder induced by  $Z_+$ .*

*Assume that  $\text{epi } h = \{(x, z) \in X \times Z \mid x \in \text{dom } h, z - h(x) \in Z_+\}$  is a lower CS-closed set of  $X \times Z$  and that  $g$  is lower CS-closed.*

*Then the composite function defined by*

$$g \circ h(x) = \begin{cases} g(h(x)) & \text{if } x \in \text{dom } h \\ +\infty & \text{otherwise} \end{cases}$$

*is lower CS-closed.*

*Proof.* Since  $g$  is nondecreasing on  $h(\text{dom } h) + Z_+$ , one has

$$g \circ h(x) = \inf_{z \in Z} g(z) + \delta_{\text{epi } h}(x, z)$$

for all  $x \in X$ . Then, by Lemma 2.1 and Corollary 2.7, the function  $(x, z) \rightarrow g(z) + \delta_{\text{epi } h}(x, z)$  is lower CS-closed. Thus  $g \circ h$  is lower CS-closed by Corollary 2.6. ■

The infimal convolution (also called epigraphical sum) and the level sum of two lower CS-closed functions are still lower CS-closed.

**COROLLARY 2.10.** *Let  $f: X \rightarrow \overline{\mathbb{R}}$  and  $g: X \rightarrow \overline{\mathbb{R}}$  be two lower CS-closed functions defined on a Fréchet space  $X$ . Then the infimal convolution of  $f$  and  $g$ ,  $f \square g$ , defined by*

$$f \square g(x) = \inf_{u \in X} f(x - u) + g(u),$$

*and the level sum of  $f$  and  $g$ ,  $f \triangle g$ , defined by*

$$f \triangle g(x) = \inf_{u \in X} f(x - u) \vee g(u),$$

*where  $a \vee b$  denotes the maximum of any extended real numbers  $a$  and  $b$ , are lower CS-closed.*

*Proof.* Set  $f_1(x, u) = f(x - u)$ ,  $g_1(x, u) = g(u)$  for all  $(x, u) \in X \times X$ . Then, by Corollary 2.9, these functions are lower CS-closed and so is the function  $f_1 + g_1$ . Now we can conclude by applying the Corollary 2.6 to the



function

$$f \sqcap g(x) = \inf_{u \in X} f_1(x, u) + g_1(x, u).$$

The proof is similar for the level sum of two functions. ■

## 2.6. Openness of the Lower CS-Closed Sets

As said in the Introduction, we are interested in the property

$$\text{core}(C) = \text{int}(C) \quad (5)$$

where  $C$  is a convex set of a l.c.s. space  $X$ .

Recall that a barrelled topological vector space is a locally convex topological vector space in which every barrel (an absorbing, convex, closed, and balanced subset) is a neighborhood of the origin.

It is known that (5) is satisfied for closed convex sets of barrelled spaces. It follows that convex semiclosed sets ( $C$  is semiclosed if  $\text{int}(\text{cl}(C)) = \text{int}(C)$ ) of barrelled spaces still satisfy (5). Moreover, it is also known (see [10, 17.B; 11]) that the CS-closed sets of metrizable topological vector spaces are semiclosed; therefore any CS-closed set of a metrizable barrelled space satisfies (5).

The next theorem says that property (5) is true for lower CS-closed sets. This is a consequence of an open mapping theorem of Rodriguez and Simons [23].

**THEOREM 2.4.** *Let  $X$  be a metrizable barrelled space. Let  $C$  be a lower CS-closed set of  $X$ . Then  $\text{core}(C) = \text{int}(C)$ .*

## 2.7. Open Mapping and Lower CS-Closed Graph Theorems

Let us first recall some definitions:

**DEFINITION 2.3.** Let  $X$  and  $Y$  be two l.c.s. spaces, let  $M: X \rightarrow Y$  be a set-valued mapping, and let  $(x, y) \in M$ .

$M$  is said to be open at  $(x, y)$  if, given any neighborhood  $U$  of  $x$ ,  $M(U)$  is a neighborhood of  $y$ .

$M$  is said to be lower semicontinuous at  $(x, y)$  if, given any neighborhood  $V$  of  $y$ ,  $M^{-1}(V)$  is a neighborhood of  $x$ .

From Theorem 2.4, one can state the following result:

**THEOREM 2.5.** *Let  $X$  be a Fréchet space, let  $Y$  be a metrizable barrelled space, let  $M: X \rightarrow Y$  be a set-valued mapping with lower CS-closed graph, and let  $(x, y) \in M$ . Suppose that  $y \in \text{core}(M(X))$ . Then  $M$  is open at  $(x, y)$ .*

We will need the following lemma:

**LEMMA 2.2** [13, 1.2.8]. *Let  $X$  be a convex subset of a product of vector spaces  $X \times Y$ . Suppose that  $(x, y) \in A$  and  $B$  is a subset of  $Y$  such that  $y \in \text{core}(B)$ . Then  $\text{core}(A_X) \subset \text{core}([A \cap (X \times B)]_X)$ .*

*Proof of Theorem 2.5.* Let  $V$  be a neighborhood of  $x$  and let  $U$  be a closed neighborhood of  $x$  contained in  $V$ . Then  $x \in \text{core}(U)$ . Using the fact that  $y \in \text{core}(M(X))$ , one can derive from Lemma 2.2 that  $y \in \text{core}(M(U))$ . On the other hand  $M(U)$  is lower CS-closed so  $y \in \text{int}(M(U)) \subset \text{int}(M(V))$ , which completes the proof. ■

**Remark 2.1.** 1. In the above theorem, if  $X$  is metrizable barrelled,  $Y$  is Fréchet, and  $x \in \text{core}(M^{-1}(Y))$ , then  $M$  is lower semicontinuous at  $(x, y)$ .

2. The preceding theorem generalizes Theorem 1 of [20]. It fits in the same line as the results of Borwein [5], Jameson [11], Kusraev and Kutateladze [13], Rodriguez and Simons [23], and Ursescu [26].

We can derive directly from the previous theorem an interesting result in the linear case:

**COROLLARY 2.11.** *Let  $X$  be a metrizable barrelled space, let  $Y$  be a Fréchet space and let  $A: X \rightarrow Y$  be a linear mapping with lower CS-closed graph. Then  $A$  is continuous.*

**Remark 2.2.** This corollary fits in the line of the results of Martineau and Schwartz on the borelian graph theorem (see [16, 17, 24]).

## 2.8. Continuity and Subdifferentiability of the Lower CS-Closed Functions

We can establish a result about the continuity of a lower CS-closed function. The following result generalizes a result which says that any finite valued lower semicontinuous convex function on a barrelled space is continuous on this space. We will need the following well-known result (see [9] for example):

**PROPOSITION 2.2.** *Let  $X$  be a l.c.s. space, let  $f: X \rightarrow \overline{\mathbb{R}}$  be a convex function, and let  $z \in \text{dom } f$ . Then  $f$  is continuous at  $z$  if and only if  $f$  is bounded above in a neighborhood of  $z$ .*

**THEOREM 2.6.** *Let  $X$  be a metrizable barrelled space and let  $f: X \rightarrow \overline{\mathbb{R}}$ . Suppose that for all  $r \in \mathbb{R}$ ,  $[f \leq r] = \{x \in X \mid f(x) \leq r\}$  is a lower CS-closed set. Then  $f$  is continuous at each point of  $\text{core}(\text{dom } f)$  which is equal to  $\text{int}(\text{dom } f)$ .*

*Proof.* If  $z \in \text{core}(\text{dom } f)$  and  $r > f(z)$  then  $z \in \text{core}([f \leq r])$  but  $[f \leq r]$  is lower CS-closed so, using Theorem 2.4, it is a neighborhood of  $z$  and, due to Proposition 2.2,  $f$  is continuous at  $z$ . ■

One of the applications of the continuity of a convex function at a point is its subdifferentiability at this point. We put this result in a more precise form. Recall that if  $f: X \rightarrow \mathbb{R}$  is a convex function over a topological vector space  $X$  and  $z$  is such that  $f(z) \in \mathbb{R}$  then the subdifferential of  $f$  at  $z$ , denoted by  $\partial f(z)$ , is the following subset of the topological dual of  $X$

$$\partial f(z) = \{x^* \in X^*: \langle x^*, x - z \rangle + f(z) \leq f(x), \forall x \in X\}.$$

We say that  $f$  is subdifferentiable at  $z$  if  $\partial f(z) \neq \emptyset$ .

**COROLLARY 2.12.** *Let  $f: X \rightarrow \overline{\mathbb{R}}$  be a lower CS-closed function defined on a l.c.s. space  $X$ . Let  $z \in X$  such that  $f(z) \in \mathbb{R}$  and*

$$\mathbb{R}^+[\text{dom } f - z] \text{ is a metrizable barrelled space.} \quad (6)$$

*Then  $f$  is subdifferentiable at  $z$ .*

*Proof.* Set  $M = \mathbb{R}^+[\text{dom } f - z]$  and  $\varphi(x) = f(x + z)$  so that  $\text{dom } \varphi = \text{dom } f - z$ ,  $\text{epi } \varphi = \text{epi } f - (z, 0)$  and  $\partial f(z) = \partial \varphi(0)$ .

$0 \in \text{core}(\text{dom } \varphi|_M)$  and  $\text{epi } \varphi|_M = \text{epi } \varphi$  is a lower CS-closed set so  $\varphi|_M$  is continuous at the origin.

There exists (due to the Hahn–Banach theorem)  $x^* \in \partial \varphi|_M(0)$ . Now another application of the Hahn–Banach theorem implies the existence of  $y^* \in Y^*$  such that  $y^*|_M = x^*$  ( $X$  is locally convex). Using the fact that  $\text{dom } \varphi = \text{dom } \varphi|_M$ , we have  $y^* \in \partial \varphi(0)$  so  $y^* \in \partial f(z)$ . ■

*Remark 2.3.* The condition (6) of the preceding corollary is very close to the Attouch–Brezis hypothesis (see Attouch and Brezis [1]).

### 3. APPLICATIONS

#### 3.1. The Use of Lower CS-Closed Functions in Convex Duality Theory

Let us recall in a few words the setting of perturbational convex duality theory in the line of Rockafellar [22]. Let  $U$  and  $X$  be two Fréchet spaces, let  $U^*$  and  $X^*$  respectively be their topological dual spaces, let  $\Phi: U \times X \rightarrow \overline{\mathbb{R}}$  be a given proper convex and lower semicontinuous function, and let  $\bar{x}^* \in X^*$ .

Consider the optimization problem

$$\text{minimize } \Phi(0, x) - \langle \bar{x}^*, x \rangle, \quad x \in X. \quad (\mathcal{P})$$

The dual problem  $(\mathcal{Q})$  associated with the primal problem  $(\mathcal{P})$  is traditionally defined as

$$\text{maximize } -\Phi^*(u^*, \bar{x}^*), \quad u^* \in U^*. \quad (\mathcal{Q})$$

Here  $\Phi^*$  is the Legendre–Fenchel conjugate function of  $\Phi$  defined on the product space  $U^* \times X^*$  by

$$\Phi^*(u^*, x^*) = \sup_{(u, x) \in U \times X} \langle u^*, u \rangle + \langle x^*, x \rangle - \Phi(u, x)$$

for each  $(u^*, x^*) \in U^* \times X^*$ .

One says that the problem  $(\mathcal{P})$  is stable if both problems  $(\mathcal{P})$  and  $(\mathcal{Q})$  have the same value, possibly  $-\infty$ , and if  $(\mathcal{Q})$  admits an optimal solution. Indeed this is the case when the following condition is fulfilled (see, for instance, [25, Theorem 2.1] for Banach spaces):

(Q.C.1)  $\mathbb{R}_+[\text{dom } \Phi]_U$  is a closed vector subspace of  $U$ .

It is important to point out that the condition (Q.C.1) does not depend on  $\bar{x}^*$  (see [4]). This result, which derives systematically from a slight extension of the Attouch–Brezis theorem on the conjugate of the sum of two convex proper lower semicontinuous functions, supposes essentially that the perturbational function  $\Phi$  is lower semicontinuous.

However, as we shall see later on, in many cases the function  $\Phi$  is *not* lower semicontinuous but lower CS-closed!

This observation is at the origin of the following result:

**THEOREM 3.1.** *Let  $\Phi: U \times X \rightarrow \overline{\mathbb{R}}$  be a lower CS-closed function. Suppose that (Q.C.1) holds. Then*

$$-\infty \leq \max_{u^* \in U^*} -\Phi^*(u^*, x^*) = \inf_{x \in X} \Phi(0, x) - \langle x^*, x \rangle < +\infty \quad (7)$$

for all  $x^* \in X^*$ . Moreover,

$$\inf_{x \in X} \Phi(0, x) - \langle x^*, x \rangle = \inf_{x \in X} \Phi^{**}(0, x) - \langle x^*, x \rangle. \quad (8)$$

*Proof.* Set

$$p(u) = \inf_{x \in X} \Phi(u, x) - \langle x^*, x \rangle$$

for  $u \in U$ . Thus  $\text{dom } p = [\text{dom } \Phi]_U$  and (Q.C.1) entails

$$\inf(\mathcal{P}) = p(0) < +\infty.$$

Moreover, one always has

$$p(0) \geq \sup_{u^* \in U^*} -\Phi^*(u^*, x^*).$$

So if  $p(0) = -\infty$ , (7) is proved.

In the following we suppose that  $p(0) \in \mathbb{R}$ . From Corollary 2.6,  $p$  is lower CS-closed. Using the condition (Q.C.1) and Corollary 2.12, we obtain that  $p$  is subdifferentiable at the origin. Therefore, there exists  $\bar{u}^* \in U^*$  such that for all  $u \in U$ ,

$$p(0) = \inf_{x \in X} \Phi(0, x) - \langle x^*, x \rangle \leq \inf_{x \in X} (\Phi(u, x) - \langle x^*, x \rangle) - \langle \bar{u}^*, u \rangle$$

so

$$\inf_{(x, u) \in X \times U} \Phi(u, x) - \langle x^*, x \rangle - \langle \bar{u}^*, u \rangle \geq \inf_{x \in X} \Phi(0, x) - \langle x^*, x \rangle.$$

As the opposite inequality obviously holds we get

$$-\Phi^*(\bar{u}^*, x^*) = \inf_{x \in X} \Phi(0, x) - \langle x^*, x \rangle,$$

which completes the proof of (7).

To prove (8) it suffices to observe that

$$\sup_{u^* \in U^*} -\Phi^*(u^*, x^*) = \sup_{u^* \in U^*} -\Phi^{***}(u^*, x^*) \leq \inf_{x \in X} \Phi^{**}(0, x) - \langle x^*, x \rangle$$

and

$$\inf_{x \in X} \Phi^{**}(0, x) - \langle x^*, x \rangle \leq \inf_{x \in X} \Phi(0, x) - \langle x^*, x \rangle = \max_{u^* \in U^*} -\Phi^*(u^*, x^*).$$

■

As a corollary, let us quote another general duality result involving marginal lower CS-closed functions:

**COROLLARY 3.1.** *Let  $\Phi: U \times X \rightarrow \bar{\mathbb{R}}$  be a lower CS-closed function. Suppose that (Q.C.1) holds. Then the convex function*

$$n: x^* \in X^* \mapsto n(x^*) = \inf_{u^* \in U^*} \Phi^*(u^*, x^*)$$

*is weak\* lower semicontinuous, exact (i.e., the infimum above is reached) and does not take the value  $-\infty$ .*

*Proof.* By Theorem 3.1,  $n$  is exact and for all  $x^* \in X^*$  one has

$$n(x^*) = - \inf_{x \in X} \Phi(0, x) - \langle x^*, x \rangle.$$

In other words

$$n(x^*) = (\Phi(0, \cdot))^*(x^*)$$

for all  $x^* \in X^*$ . Consequently  $n$  is weak\* lower semicontinuous.

Let us prove that  $n$  does not take the value  $-\infty$ . Suppose the contrary. Then

$$(\Phi(0, \cdot))^* = n^*(\cdot) = +\infty,$$

and, a fortiori,

$$\Phi(0, x) = +\infty \quad \forall x \in X.$$

Hence  $0 \notin [\text{dom } \Phi]_U$ , a contradiction with (Q.C.1). ■

The corollary below extends the Rockafellar and Attouch-Brezis theorems (see also [23]) to lower CS-closed functions:

**COROLLARY 3.2.** *Let  $f, g: X \rightarrow \overline{\mathbb{R}}$  be two lower CS-closed functions on the Fréchet space  $X$  such that*

$$\mathbb{R}^+ [\text{dom } f - \text{dom } g] \text{ is a closed linear subspace of } X.$$

*Then we have  $(f + g)^* = f^* \square g^*$ . In particular, the inf-convolution  $f^* \square g^*$  is convex and weak\* lower semicontinuous. Moreover,  $f^* \square g^*$  is exact (at each point of  $X^*$ ) and does not take the value  $-\infty$ .*

*Proof.* For  $(u, x) \in X \times X$  let

$$\Phi(u, x) = f(u - x) + g(x).$$

Then, by Corollaries 2.9 and 2.7,  $\Phi$  is lower CS-closed and

$$[\text{dom } \Phi]_X = \text{dom } g - \text{dom } f.$$

On the other hand the Legendre–Fenchel conjugate of  $\Phi$  is

$$\Phi^*(u^*, x^*) = f^*(-u^*) + g^*(u^* + x^*)$$

for all  $(u^*, x^*) \in X^* \times X^*$ . By Theorem 3.1, one thus has

$$n(\cdot) = \inf_{u^* \in X^*} \Phi^*(u^*, \cdot) = (f^* \square g^*)(\cdot).$$

Moreover  $n$  is exact and

$$n(x^*) = - \inf_{x \in X} \Phi(0, x) - \langle x^*, x \rangle = (f + g)^*(x^*)$$

for all  $x^* \in X^*$ , so that the proof is complete. ■

### 3.2. On the Conjugate of the Sum of a Lower CS-Closed Function and a Lower CS-Closed Composite Function

Assume that the Fréchet space  $Z$  is equipped with a preorder relation induced by a convex cone  $Z_+$  of  $Z$ : for  $z_1, z_2 \in Z$  we set

$$z_1 \leq z_2 \quad \text{if } z_2 - z_1 \in Z_+.$$

Let  $h$  be a function defined on a nonempty convex subset  $\text{dom } h$  of another Fréchet space  $X$  and taking its values in  $Z$ . We denote by

$$\text{epi } h = \{(x, z) \in \text{dom } h \times Z \mid h(x) \leq z\}$$

the epigraph of  $h$  and assume that  $\text{epi } h$  is convex.

Let us consider two extended-real-valued convex functions,  $f: X \rightarrow \overline{\mathbb{R}}$ ,  $g: Z \rightarrow \overline{\mathbb{R}}$ , with  $g$  nondecreasing on the subset  $h(\text{dom } h) + Z_+$  of  $Z$ . Assuming that  $\text{epi } h$  is closed,  $f \in \Gamma_0(X)$ , and  $g \in \Gamma_0(Z)$ , the expression of the conjugate function of the convex function  $f + g \circ h$  has been explicated in [7] under the following condition:

(Q.C.2)

$\mathbb{R}_+[\text{dom } g - h(\text{dom } f \cap \text{dom } g) - Z_+]$  is a closed vector subspace of  $Z$ .

By using Theorem 3.1 we intend to recapture and generalize this result assuming only the set  $\text{epi } h$  and the functions  $f, g$  are lower CS-closed (see [19] for another generalization).

Let us observe that the convex function  $f + g \circ h$  defined on  $X$  by

$$f + g \circ h(x) = \begin{cases} f(x) + g(h(x)) & \text{if } x \in \text{dom } h, \\ +\infty & \text{otherwise} \end{cases}$$

may be written under the form

$$f + g \circ h(x) = \Phi(0, x)$$

where  $\Phi$  is defined on  $Z \times X$  by

$$\Phi(u, x) = \begin{cases} \inf_{z \in Z} f(x) + g(z) + \delta_{\text{epi } h}(x, z - u) & \text{if } x \in \text{dom } h, \\ +\infty & \text{otherwise.} \end{cases}$$

Now observe that the conjugate function of  $f + g \circ h$  is given by

$$(f + g \circ h)^*(x^*) = - \inf_{x \in X} \Phi(0, x) - \langle x^*, x \rangle$$

for each  $x^* \in X^*$ .

Here again the function  $\Phi$  is *not necessarily lower semicontinuous* but it is not difficult to show that  $\Phi$  is lower CS-closed. Now observing that

$$[\text{dom } \Phi]_U = \text{dom } g - h(\text{dom } h \cap \text{dom } f) - Z_+,$$

one sees that condition (Q.C.1) is satisfied.

To apply Theorem 3.1 we need the conjugate function of  $\Phi$ . To this end, let us introduce the positive cone  $Z_+^*$  of  $Z^*$  consisting of all the continu-

ous positive linear forms  $u^* \in Z_+^*$ , that is,

$$\langle u^*, u \rangle \geq 0, \quad \forall u \in Z_+.$$

In what follows, we extend the functions  $u^* \circ h$  to the whole space  $X$  by setting  $(u^* \circ h)(x) = +\infty$  if  $x \notin \text{dom } h$ . Hence we have the following result:

LEMMA 3.1. *For each  $(u^*, x^*) \in Z^* \times X^*$ , we have*

$$\Phi^*(u^*, x^*) = \begin{cases} (f + u^* \circ g)(x^*) + g^*(u^*) & \text{if } u^* \in Z_+^*, \\ +\infty & \text{otherwise.} \end{cases}$$

Now, we derive the expression of the conjugate function of  $f + g \circ h$ , generalizing [7, Proposition 4.11]:

COROLLARY 3.3. *Let  $X, Z$  be two Fréchet spaces,  $Z$  being equipped with a partial order induced by a convex cone  $Z_+$ , let  $f$  and  $g$  be two lower CS-closed functions on  $X$  and  $Z$  respectively, and let  $h$  be a mapping defined on a convex subset  $\text{dom } h$  of  $X$  with values in  $Z$  and such that  $\text{epi } h$  is lower CS-closed. Assume that  $g$  is nondecreasing on the subset  $h(\text{dom } h) + Z_+$  and that (Q.C.2) holds. Then we have*

$$(f + g \circ h)^*(x^*) = \min_{u^* \in Z_+^*} g^*(u^*) + (f + u^* \circ h)^*(x^*)$$

for each  $x^* \in X^*$ .

*Proof.* Applying Theorem 3.1 to the function  $\Phi$ , we can get the announced result. ■

### 3.3. Asymptotic calculus

Let  $U, X$  be two Fréchet spaces,  $G$  a convex proper weak\* lower semicontinuous function on  $U^* \times X^*$ , and  $g(x^*) = \inf_{u^* \in U^*} G(u^*, x^*)$ . Suppose that

(Q.C.3)  $\mathbb{R}_+[\text{dom } G^*]_U$  is a closed vector subspace of  $U$ ,

where the conjugate function  $G^*$  of  $G$  is considered on the product space  $U \times X$  and not  $U^{**} \times X^{**}$ .

By Corollary 3.1 apply to  $\Phi = G^*$ ,  $g$  is convex weak\* lower semicontinuous, exact, and does not take the value  $-\infty$ . Furthermore  $g^*(\cdot) = G^*(0, \cdot)$  so that,  $G^*$  being proper,  $g$  is not identically  $+\infty$ .

It follows that the asymptotic functional  $g_\infty$  of  $g$  coincides with the support function of the effective domain of  $g^*$ , i.e., for each  $x^* \in X^*$ , we have (see [14])

$$g_\infty(x^*) = \sup_{x \in \text{dom } g^*} \langle x^*, x \rangle.$$



On the other hand

$$g^*(x) = G^*(0, x), \quad \forall x \in X;$$

we then have

$$g_\infty(x) = \sup \langle x^*, x \rangle \mid G^*(0, x) < +\infty,$$

or, equivalently,

$$-g_\infty(x) = \inf_{x \in X} \delta_{\text{dom } G^*}(0, x) - \langle x^*, x \rangle.$$

Of course  $\delta_{\text{dom } G^*}$  is not necessarily lower semicontinuous nor CS-closed. But  $\delta_{\text{dom } G^*}$  is actually lower CS-closed. Now we can state:

**THEOREM 3.2.** *Let  $G \in \Gamma_0(U^* \times X^*)$  and let  $g: X^* \rightarrow \overline{\mathbb{R}}$  be defined by*

$$g(x^*) = \inf_{u^* \in U^*} G(u^*, x^*).$$

*Suppose that (Q.C.3) holds. Then  $g$  is convex, proper, weak\* l.s.c., exact and one has*

$$g_\infty(x^*) = \min_{u^* \in U^*} G_\infty(u^*, x^*) \quad (9)$$

*for all  $x^* \in X^*$ .*

*Proof.* We only need to prove (9).

Applying Theorem 3.1 to  $\Phi = \delta_{\text{dom } G^*}$ , we then have

$$\max_{u^* \in U^*} -\delta_{\text{dom } G^*}^*(u^*, x^*) = \inf_{x \in X} \delta_{\text{dom } G^*}(0, x) - \langle x^*, x \rangle = -g_\infty(x) < +\infty$$

but  $\delta_{\text{dom } G^*}^* = G_\infty$  and the proof is complete. ■

**Remark 3.1.** The formula (9) was established (see [29]) in the setting of locally convex space under some relative compactness assumptions which we do not need here (see also [3]).

As an example, let us consider the inf-convolution of two convex proper weak\* lower semicontinuous functions  $\varphi$  and  $\psi$  defined on  $X^*$ . We know that if

(Q.C.4)

$$\mathbb{R}_+(\text{dom } \varphi^* - \text{dom } \psi^*) \text{ is a closed vector subspace of } X$$

then the inf-convolution  $\varphi \square \psi$  defined by

$$x^* \in X^* \rightarrow (\varphi \square \psi)(x^*) = \inf_{u^* \in U^*} \varphi(x^* - u^*) + \psi(u^*)$$

is a convex proper weak\* lower semicontinuous function (see [1]).

The formula

$$(\varphi \square \psi)_\infty = \varphi_\infty \square \psi_\infty$$

was established in the hilbertian setting in [2], by means of the asymptotic function associated with a monotone maximal operator, under the hypothesis

$$\text{int}(\text{dom } \partial\varphi^*) \cap \text{dom } \partial\psi^* \neq \emptyset.$$

In fact we have the following result which allows us to weaken sensibly the above condition

**COROLLARY 3.4.** *Let  $\varphi, \psi \in \Gamma_0(X^*)$ . Suppose that (Q.C.4) holds. Then*

$$(\varphi \square \psi)_\infty = \varphi_\infty \square \psi_\infty.$$

*Proof.* Let

$$G(u^*, x^*) := \varphi(x^* - u^*) + \psi(u^*)$$

for all  $u^*, x^*$  in  $X^*$ . Then we have

$$G^*(u, x) = \varphi^*(x) + \psi^*(u + x)$$

for all  $u, x$  in  $X$ . It follows that

$$[\text{dom } G^*]_U = \text{dom } \psi^* - \text{dom } \varphi^* \quad \text{and}$$

$$G_\infty(u^*, x^*) = \varphi_\infty(x^* - u^*) + \psi_\infty(u^*).$$

From Theorem 3.2 we obtain the announced result. ■

## ACKNOWLEDGMENTS

The authors are grateful to Professors J. P. Penot, L. Thibault, M. Valadier, and M. Volle for their hints and insightful comments.

## REFERENCES

1. H. Attouch and H. Brezis, Duality for the sum of convex functions in general Banach spaces, in "Aspects of Mathematics and Its Applications" (J. A. Barroso, Ed.), Elsevier, Amsterdam/New York, 1986.
2. H. Attouch, Z. Chbani, and A. Moudafi, Recession operator and solvability of noncoercive optimization and variational problem, *Sém. Anal. Convexe* **5** (1994).
3. A. Auslender, Non coercive optimization problems, *Math. Oper. Res.* **21**, No. 4 (1996), 769–789.

4. A. Auslender, R. Cominetti, and J. P. Crouzeix, Convex functions with unbounded level sets and applications to duality theory, *SIAM J. Optim.* **3**, No. 4 (1993), 669–687.
5. J. M. Borwein, Convex relations in analysis and optimization, in “Generalized Concavity in Optimization and Economics,” (S. Schaible and T. Ziemmba, Eds.), Academic Press, San Diego, 1981.
6. N. Bourbaki, “Espaces vectoriels topologiques,” Chap. 1–5, Masson, Paris, 1981.
7. C. Combari, M. Laghdir, and L. Thibault, Sous-différentiels de fonctions composées, *Ann. Sci. Math. Québec* **18**, No. 2 (1994), 119–148.
8. R. Cominetti, Some remarks on convex duality in normed spaces with and without compactness, *Control Cybernet.* **23**, No. 1/2 (1994), 123–138.
9. Y. Ekeland and R. Temam, “Analyse convexe et problèmes variationnels,” Dunod, Paris, 1974.
10. R. B. Holmes, “Geometric Functional Analysis and Its Applications,” Springer-Verlag, Berlin/New York, 1975.
11. J. O. Jameson, Convex series, *Proc. Cambridge Philos. Soc.* **72** (1972), 37–47.
12. A. G. Kusraev, On openness of measurable convex correspondences, *Mat. Zametki* **1** (1983), 41–48.
13. A. G. Kusraev and S. S. Kutateladze, “Subdifferentials: Theory and Applications,” Mathematics and Its Applications, Kluwer Academic, Dordrecht/Norwell, MA, 1992.
14. J. P. Laurent, “Approximation et Optimisation,” Hermann, Paris, 1972.
15. E. A. Lifshits, Ideally convex sets, *Funct. Anal. Appl.* **4**, No. 4 (1970), 76–77.
16. A. Martineau, Sur le théorème du graphe fermé, *C. R. Acad. Sci. Paris Sér. A* **263** (1966), 870–871.
17. A. Martineau, Sur des théorèmes de S. Banach et de L. Schwartz concernant le graphe fermé, *Studia Math.* **30** (1968), 43–51.
18. J. J. Moreau, “Fonctionnelles convexes,” Collège de France, 1966.
19. M. Moussaoui and M. Volle, Quasicontinuity and united functions in convex duality theory, *Comm. Appl. Nonlinear Anal.* **4** (1997), 73–89.
20. S. M. Robinson, Regularity and stability for convex multivalued functions, *Math. Oper. Res.* **1** (1976), 130–143.
21. R. T. Rockafellar, “Convex Analysis,” Princeton Univ. Press, Princeton, NJ, 1970.
22. R. T. Rockafellar, Conjugate duality and optimization, in SIAM Regional Conference in Applied Mathematics, Vol. 16, SIAM, Philadelphia, 1974.
23. B. Rodriguez and S. Simons, Conjugate functions and subdifferentials in non-normed situations for operators with complete graphs, *Nonlinear Anal.* **12** (1988), 1062–1078.
24. L. Schwartz, Sur le théorème du graphe fermé, *C. R. Acad. Sci. Paris Sér. A* **263** (1966), 602–605.
25. S. Traoré and M. Volle, On the level sum of two convex functions on Banach spaces, *J. Convex Anal.* **3**, No. 1 (1996), 141–151.
26. C. Ursescu, Multifunctions with convex closed graph, *Czechoslovak Math. J.* **25** (1975), 438–441.
27. M. Volle, Sur quelques formules de dualité convexe et non convexe, *Set-Valued Anal.* **2** (1994), 369–379.
28. C. Zalinescu, On some open problems in convex analysis, *Arch. Math.* **59** (1992), 566–571.
29. C. Zalinescu, Stability for a class of non linear optimization problems and applications, in “Nonsmooth Optimization and Related Topics,” Plenum, New York, 1989.